

A UNIFORM KADEC-KLEE PROPERTY FOR SYMMETRIC OPERATOR SPACES

P.G. Dodds, T.K. Dodds, P.N. Dowling, C.J. Lennard*† and F.A. Sukochev*

ABSTRACT

We show that if a rearrangement invariant Banach function space E on the positive semi-axis satisfies a non-trivial lower q -estimate with constant 1 then the corresponding space $E(\mathcal{M})$ of τ -measurable operators, affiliated with an arbitrary semi-finite von Neumann algebra $\mathcal{M}$ equipped with a distinguished faithful, normal, semi-finite trace τ , has the uniform Kadec-Klee property for the topology of local convergence in measure. In particular, the Lorentz function spaces $L_{q,p}$ and the Lorentz-Schatten classes $\mathcal{C}_{q,p}$ have the UKK property for convergence locally in measure and for the weak-operator topology, respectively. As a partial converse, we show that if E has the UKK property with respect to local convergence in measure then E must satisfy some non-trivial lower q -estimate. We also prove a uniform Kadec-Klee result for local convergence in any Banach lattice satisfying a lower q -estimate.

1. Introduction and preliminaries

It is well-known (see, for example, [Ar],[Si]) that the Schatten p -classes $\mathcal{C}_p$, $1 \leq p < \infty$, have the (so-called) Kadec-Klee (or Radon-Riesz) property, that is norm convergence in $\mathcal{C}_p$ coincides with weak operator convergence for sequences in the unit sphere. It has been shown recently by Lennard [Le1] that the direct argument given by Arazy [Ar] for trace-class operators may be refined to show that the Schatten class $\mathcal{C}_1$ has the uniform

* Research partially supported by A.R.C.

† Research partially supported by a U. Pittsburgh FAS Grant

1980 *Mathematics Subject Classification*. Primary 46E30; Secondary 46L50, 47B55.

Kadec-Klee property for the weak operator topology (see below for the precise definitions). Further extensions to the setting of trace ideals have been announced by Sukochev [Su]. From a somewhat different (though related) viewpoint, it is a classical theorem of F. Riesz that norm convergence for sequences in the unit sphere of $L^1([0, 1])$ coincides with convergence in measure. Appropriate uniform versions of this theorem may be found in [Le2], and extensions to certain Lorentz spaces in [Le3].

It is the intention of the present paper to give simultaneous extensions of each of the above theorems in the setting of symmetric spaces of measurable operators. Our approach is based on a notion of local convergence in measure for measurable operators which specialises to the weak operator topology for the case of trace ideals in the sense of Gohberg and Krein [GK], to pointwise convergence in the case of symmetric sequence spaces, and to convergence in measure on sets of finite measure in the case of rearrangement invariant function spaces. Let $\mathcal{M}$ be a semi-finite von Neumann algebra, equipped with a distinguished faithful, normal semi-finite trace τ . Our principal result (Theorem 2.7) shows that if E is a symmetric (rearrangement-invariant) quasi-Banach function space on the semi-axis $\mathbf{R}^+$ which satisfies a non-trivial lower estimate with constant 1, then the corresponding symmetric operator space $E(\mathcal{M})$ has the uniform Kadec-Klee property for local convergence in measure. Moreover, our theorem is best possible in the sense that symmetric spaces on the positive half-line which have this uniform property must necessarily satisfy a non-trivial lower estimate. The existence of such an estimate is a geometric condition which is readily verified, and so our results essentially give a useful geometric characterisation of this uniform Kadec-Klee type property which not only unifies the results alluded to above, but also yields an extensive class of new examples. In each of these examples, our principal theorem yields results concerning normal structure and fixed point properties. For a more detailed discussion of these aspects and the relation between normal structure and uniform Kadec-Klee type properties, we refer the interested reader

to [Le2], and the references contained therein.

We remark that Dilworth and Hsu [DH] have recently proved a uniform version of the theorem of [Ar]/[Si] i.e. they show that if a symmetric sequence space E has the UKK property for pointwise convergence, then the associated unitary matrix space $\mathcal{C}_E$ has the UKK property for the weak operator topology. They complement this Banach space result by showing that the Schatten class $\mathcal{C}_p$ has UKK(weak operator) for all $0 < p \leq 1$, which is a special case of our Theorem 2.7 below. Along the way they prove a special case of our main inequality in Proposition 2.5. However, their methods are based on a generalization of McCarthy's characterization of Schatten class norms in [M], rather than inequalities for decreasing rearrangements of operators; and so differ from the methods we have used.

We begin by recalling some necessary terminology. Let E be a complex quasi-Banach lattice. If $0 < p < \infty$, then E is said to be p -convex (resp. p -concave) if there exists a constant $C > 0$ such that for all finite sequences (x_n) in E

$$\begin{aligned} \|(\sum |x_n|^p)^{\frac{1}{p}}\|_E &\leq C(\sum \|x_n\|_E^p)^{\frac{1}{p}}, \\ (\text{resp. } C\|(\sum |x_n|^p)^{\frac{1}{p}}\|_E &\geq (\sum \|x_n\|_E^p)^{\frac{1}{p}}). \end{aligned}$$

The least such constant C is called the p -convexity (resp. p -concavity) constant of E and is denoted by $M^{(p)}(E)$ (resp. $M_{(p)}(E)$). For $0 < p < \infty$, $E^{(p)}$ will denote the quasi-Banach lattice defined by

$$E^{(p)} = \{x : |x|^p \in E\}$$

equipped with the quasi-norm

$$\|x\|_{E^{(p)}} = \| |x|^p \|_E^{1/p}.$$

It is not difficult to see that if E is α -convex and q -concave then $E^{(p)}$ is αp -convex and qp -concave with $M^{(\alpha p)}(E^{(p)}) \leq M^{(\alpha)}(E)^{\frac{1}{p}}$ and $M_{(qp)}(E^{(p)}) \leq M_{(q)}(E)^{\frac{1}{p}}$. Conse-

quently, if E is α -convex then $E^{(1/\alpha)}$ is 1-convex and so can be renormed as a Banach lattice (cf.[LT]).

The quasi-Banach lattice E is said to satisfy a *lower q -estimate* (resp. *upper p -estimate*) if there exists a positive constant $C > 0$ such that for all finite sequences (x_n) of mutually disjoint elements in E

$$\begin{aligned} (\sum \|x_n\|_E^q)^{1/q} &\leq C \|\sum x_n\|_E, \\ (\text{resp. } C(\sum \|x_n\|_E^p)^{1/p} &\geq \|\sum x_n\|_E). \end{aligned}$$

It is clear that if E is q -concave for some $q \in (0, \infty)$, then E satisfies a lower q -estimate; if E satisfies a lower q -estimate for some $q \in (1, \infty)$ then E is r -concave, for all $r > q$ [LT]. However, as shown in [Cr] Theorem 3.4(i), the Lorentz space $L_{q,p}(\mathbf{R}^+)$, $1 \leq p < q$ satisfies a lower q -estimate, but is not q -concave.

We denote by $\mathcal{M}$ a semi-finite von Neumann algebra in the Hilbert space H with given normal faithful semi-finite trace τ . If x is a (densely defined) self-adjoint operator in H and if $x = \int_{(-\infty, \infty)} s de_s^x$ is its spectral decomposition then, for any Borel subset $B \subseteq \mathbf{R}$, we denote by $\chi_B(x)$ the corresponding spectral projection $\int_{(-\infty, \infty)} \chi_B(s) de_s^x$. A closed densely defined linear operator x in H affiliated with $\mathcal{M}$ is said to be τ -measurable if and only if there exists a number $s \geq 0$ such that

$$\tau(\chi_{(s, \infty)}(|x|)) < \infty.$$

The set of all τ -measurable operators will be denoted by $\widetilde{\mathcal{M}}$. The set $\widetilde{\mathcal{M}}$ is a *-algebra with sum and product being the respective closures of the algebraic sum and product. For $x \in \widetilde{\mathcal{M}}$, the *generalized singular value function* (or *decreasing rearrangement*) $\mu_*(x)$ of x is defined by

$$\mu_t(x) = \inf\{s \geq 0 : \tau(\chi_{(s, \infty)}(|x|)) \leq t\}, \quad t \geq 0.$$

It should be noted that the decreasing rearrangement $\mu(x)$ is the right continuous, non-increasing inverse of the distribution function $\lambda_s(x)$, where $\lambda_s(x) = \tau(\chi_{(s,\infty)}(|x|))$, $s \geq 0$. It may be shown (cf. [FK]) that if $x \in \widetilde{\mathcal{M}}$ and if $t > 0$ then

$$\mu_t(x) = \inf\{\|xe\|_{\mathcal{M}} : e \in \mathcal{M} \text{ a projection with } \tau(1-e) \leq t\}.$$

It follows that $\mu(x)$ is a decreasing, right-continuous function on the half-line $\mathbf{R}^+ = [0, \infty)$, and that $\mu(x) = \mu(x^*) = \mu(|x|)$, for all $x \in \widetilde{\mathcal{M}}$. For basic properties of decreasing rearrangements of measurable operators, we refer to [FK]. The topology defined by the translation invariant metric d on $\widetilde{\mathcal{M}}$ obtained by setting

$$d(x, y) = \inf\{t \geq 0 : \mu_t(x - y) \leq t\}, \quad \text{for } x, y \in \widetilde{\mathcal{M}},$$

is called the *measure topology*. It is shown in [Ne] and [Te] that $\widetilde{\mathcal{M}}$ equipped with the measure topology is a complete, Hausdorff, topological $*$ -algebra in which $\mathcal{M}$ is dense. It is not difficult to see that a net $(x_i)_{i \in I}$ in $\widetilde{\mathcal{M}}$ converges to a measurable operator x for the measure topology if and only if for every $\epsilon > 0$, $\delta > 0$ there exists $i_0 \in I$ such that whenever $i \geq i_0$ there exists a projection $e \in \mathcal{M}$ such that

$$\|(x_i - x)e\|_{\mathcal{M}} < \epsilon \quad \text{and} \quad \tau(1 - e) < \delta.$$

We denote by $\widetilde{\mathcal{M}}_0$ the set of all $x \in \widetilde{\mathcal{M}}$ such that $\mu_t(x) \rightarrow 0$ as $t \rightarrow \infty$.

If $\mathcal{M}$ is the space $\mathcal{L}(H)$ of all bounded linear operators on H and τ is the standard trace, then $\widetilde{\mathcal{M}} = \mathcal{L}(H)$ and $x \in \mathcal{L}(H)$ is compact if and only if $\mu_t(x) \rightarrow 0$ as $t \rightarrow \infty$, in which case for each $n = 0, 1, 2, \dots$,

$$\mu_n(x) = \mu_t(x), \quad t \in [n, n+1),$$

and $\{\mu_n(x)\}_{n=0}^{\infty}$ is the usual singular value sequence of x in decreasing order, counted according to multiplicity [GK].

We identify the space $L^\infty(\mathbf{R}^+)$ of all bounded complex-valued Lebesgue measurable functions on the half-line $\mathbf{R}^+$ as a commutative von Neumann algebra acting by multiplication on the Hilbert space $L^2(\mathbf{R}^+)$ with trace given by integration with respect to Lebesgue measure. In this case the τ -measurable operators coincide with those complex measurable functions f on $\mathbf{R}^+$ which are bounded except on a set of finite measure. In this example, the generalized singular value function, which we continue to denote by $\mu(f)$, coincides with the familiar right-continuous decreasing rearrangement of the function $|f|$. See, for example, [KPS].

Following [X], by a symmetric quasi-Banach function space on the positive half-line $\mathbf{R}^+$ is meant a quasi-Banach lattice E of measurable functions with the following properties: (i) E is an order-ideal in the linear space $L^0(\mathbf{R}^+)$ of all finite almost-everywhere measurable functions on $\mathbf{R}^+$; (ii) E is rearrangement-invariant in the sense that whenever $x \in E$ and y is a measurable function with $\mu(y) = \mu(x)$, it follows that $y \in E$ and $\|y\|_E = \|x\|_E$; (iii) E contains all finitely-supported simple functions. The norm on E is said to be a σ -Fatou norm if

$$0 \leq x_n \uparrow x, \quad x_n, x \in E \text{ implies } \|x_n\|_E \uparrow \|x\|_E.$$

The norm on E is said to be *order-continuous* if

$$0 \leq x_\beta \downarrow_\beta 0 \text{ implies } \|x_\beta\|_E \downarrow_\beta 0.$$

We remark that if E has order-continuous norm, then $\lim_{t \rightarrow \infty} \mu_t(x) = 0$ for all $x \in E$. It is clear that if E has order-continuous norm, then the norm on E is a σ -Fatou norm. Further, it is not difficult to see that if E is q -concave for some $0 < q < \infty$, then the norm on E is order-continuous. Now suppose that the symmetric quasi-Banach function space E is α -convex for some $0 < \alpha < \infty$ and that $M^{(\alpha)}(E) = 1$. Let F be the Banach lattice $E^{(1/\alpha)}$. If the norm on E is a σ -Fatou norm, then so is the norm on F and consequently the natural embedding of F into the second associate space (Köthe bidual) F'' is an

isometry, by a well known result of Lorentz and Luxemburg [LT] 1.b.18. Consequently, by [KPS] II.4.6, there exists a family W of non-increasing functions on $\mathbf{R}^+$ such that

$$\|x\|_F = \sup \left\{ \int_{[0,\infty)} \mu_t(x) w(t) dt : w \in W \right\} \quad (1.1)$$

for all $x \in F$. Consequently,

$$\|x\|_E^\alpha = \sup \left\{ \int_{[0,\infty)} \mu_t^\alpha(x) w(t) dt : w \in W \right\} \quad (1.2)$$

for all $x \in E$.

Unless stated otherwise, E will always denote a symmetric quasi-Banach function space on the positive half-line $\mathbf{R}^+$ and $(\mathcal{M}, \tau)$ a semifinite von Neumann algebra on a Hilbert space H , with faithful normal semi-finite trace τ . We define the symmetric space $E(\mathcal{M})$ of measurable operators associated with E by setting

$$E(\mathcal{M}) = \{x \in \widetilde{\mathcal{M}} : \mu(x) \in E\}$$

and

$$\|x\|_{E(\mathcal{M})} = \|\mu(x)\|_E, \quad x \in E(\mathcal{M}).$$

It follows immediately that $\|x\|_{E(\mathcal{M})} = \||x|\|_{E(\mathcal{M})} = \|x^*\|_{E(\mathcal{M})}$ for all $x \in E(\mathcal{M})$. It is shown in [X] Lemma 4.1 that if E has σ -Fatou norm and is α -convex for some $0 < \alpha < \infty$ with $M^{(\alpha)}(E) = 1$, then the functional $\|\cdot\|_{E(\mathcal{M})}$ is a norm if $\alpha \geq 1$ and an α -norm if $0 < \alpha < 1$. This means that if $x, y \in E(\mathcal{M})$, then

$$\|x + y\|_{E(\mathcal{M})}^\alpha \leq \|x\|_{E(\mathcal{M})}^\alpha + \|y\|_{E(\mathcal{M})}^\alpha.$$

Equipped with this norm or α -norm, the space $E(\mathcal{M})$ is complete. For related results, see [Ov], [DDP1,2].

If $x, y \in \widetilde{\mathcal{M}}$, we say that x is *submajorized by* y , written $x \prec\prec y$, if and only if

$$\int_0^\alpha \mu_t(x) dt \leq \int_0^\alpha \mu_t(y) dt, \quad \text{for all } \alpha \geq 0.$$

If E is a symmetric Banach function space with σ -Fatou norm and if $x, y \in E$ satisfy $x \prec\prec y$, then $\|x\|_E \leq \|y\|_E$; if E has order-continuous norm or is *maximal* [KPS] in the sense that the natural embedding of E into its second associate space is an isometric surjection, then E has the following property: if $y \in E$ and if $x \prec\prec y$, then $x \in E$ and $\|x\|_E \leq \|y\|_E$. Suppose now that E is α -convex with $M^{(\alpha)}(E) = 1$ for some $\alpha \in (0, \infty)$, and that $0 < q < \infty$. If $x, y \in \widetilde{\mathcal{M}}$, it follows from [FK], Theorem 4.2(iii) that

$$\mu^{\alpha q}(xy) \prec\prec \mu^{\alpha q}(x)\mu^{\alpha q}(y).$$

It now follows from the representation (1.2), the usual Hölder inequality and a well-known rearrangement inequality of Hardy [KPS] II.2.18, that for $0 < q_0, q_1, q < \infty$ with $1/q = 1/q_0 + 1/q_1$

$$\|xy\|_{E^{(q)}(\mathcal{M})} \leq \|x\|_{E^{(q_0)}(\mathcal{M})} \|y\|_{E^{(q_1)}(\mathcal{M})}. \quad (1.3)$$

We shall need the following result, which is well known for the case that E is a Banach lattice [DDP2], [CS]. Here, order-continuity of the norm on $E(\mathcal{M})$ is defined in the obvious way.

Proposition 1.1 *Let E be α -convex with $M^{(\alpha)}(E) = 1$ for some $\alpha \in (0, 1]$.*

- (i). *If E has order-continuous norm, then so does $E(\mathcal{M})$.*
- (ii). *If E has order-continuous norm, and if $\{e_\beta\} \subseteq \mathcal{M}$ is any family of (self-adjoint) projections for which $e_\beta \downarrow_\beta 0$ then $\lim_\beta \|xe_\beta\|_{E(\mathcal{M})} = 0$ for every $x \in E(\mathcal{M})$.*

Proof. (i). Suppose that E has order-continuous norm and that $0 \leq x_\beta \downarrow_\beta 0$ holds in $E(\mathcal{M})$. As noted earlier, order-continuity of the norm on E implies that $\lim_{t \rightarrow \infty} \mu_t(x) = 0$ for all $x \in E$. It follows from [DDP2] (see also [CS]) that $\mu(x_\beta) \downarrow_\beta 0$ holds in E and (i) now follows.

(ii). Suppose that $\{e_\beta\} \subseteq \mathcal{M}$ is any family of projections in $\mathcal{M}$ with $e_\beta \downarrow_\beta 0$ and let

$0 \leq x \in \mathcal{M}$. Using (1.3) above, it follows that

$$\begin{aligned} \|xe_\beta\|_{E(\mathcal{M})} &= \|e_\beta x^{1/2} x^{1/2}\|_{E(\mathcal{M})} \\ &\leq \|(x^{1/2})^2\|_{E(\mathcal{M})}^{1/2} \| |e_\beta x^{1/2}|^2 \|_{E(\mathcal{M})}^{1/2} \\ &= \|x\|_{E(\mathcal{M})}^{1/2} \|x^{1/2} e_\beta x^{1/2}\|_{E(\mathcal{M})}^{1/2}. \end{aligned}$$

Now $e_\beta \downarrow_\beta 0$ in $\mathcal{M}$ implies that $x^{1/2} e_\beta x^{1/2} \downarrow_\beta 0$ in $E(\mathcal{M})$ and the assertion of (ii) now follows from that of (i).

2. A uniform Kadec-Klee property

It is now convenient to localize the notion of convergence in measure as follows. If $\epsilon, \delta > 0$ and if e is a projection in $\mathcal{M}$ with $\tau(e) < \infty$, then the family of all sets $N_{\epsilon, \delta, e}$ consisting of all $x \in \widetilde{\mathcal{M}}$ such that $\mu_\delta(exe) < \epsilon$ form a neighbourhood base at 0 for a Hausdorff linear topology on $\widetilde{\mathcal{M}}$. This topology (cf. [DDP2], Proposition 5.13) will be called the topology of *local convergence in measure* (*lcm*). If we observe that

$$(|exe| - \lambda 1)^+ = (|exe| - \lambda e)^+$$

for all $\lambda > 0$, $x \in \widetilde{\mathcal{M}}$ and projections $e \in \mathcal{M}$, then it follows that $\mu(exe) = \mu^e(exe)$ for all $x \in \widetilde{\mathcal{M}}$, where μ^e denotes decreasing rearrangement calculated relative to the von Neumann algebra $e\mathcal{M}e$ with respect to the trace $\tau(e \cdot e)$. This remark follows also as a special case of [Fa], Proposition 1.5(i). It follows that convergence locally in measure coincides with convergence for the measure topology relative to $(e\mathcal{M}e, \tau(e \cdot e))$, for each projection $e \in \mathcal{M}$ with $\tau(e) < \infty$.

We remark immediately that if $\mathcal{M}$ is commutative, and identified with the von Neumann algebra of all multiplication operators given by bounded measurable functions on some localisable measure space then the preceding notion of local convergence in measure reduces to the more familiar notion of convergence in measure on sets of finite measure

in the underlying measure space. On the other hand, if $\mathcal{M}$ is the space $\mathcal{L}(H)$ of all bounded linear operators on H and τ is the standard trace, then convergence locally in measure is precisely convergence for the weak operator topology.

Proposition 2.1 *Assume that E is α -convex for some $0 < \alpha \leq 1$ with $M^{(\alpha)}(E) = 1$. If E has order-continuous norm, and if $\{x_n\}$ is a sequence in $E(\mathcal{M})$ which converges to $x \in E(\mathcal{M})$ locally in measure, then for every $\epsilon > 0$, there exists a projection e in $\mathcal{M}$ with $\tau(e) < \infty$ and a subsequence $\{y_n\} \subseteq \{x_n\}$ such that*

$$\|e(y_n - x)e\|_{E(\mathcal{M})} \rightarrow_n 0 \text{ and } \max\{\|x(1 - e)\|_{E(\mathcal{M})}, \|x^*(1 - e)\|_{E(\mathcal{M})}\} < \epsilon.$$

Proof The semi-finiteness of $\mathcal{M}$ implies that the family $\{e_\beta\}$ of all projections in $\mathcal{M}$ of finite trace satisfies $0 \leq e_\beta \uparrow_\beta 1$. Since E has order-continuous norm, it follows from Proposition 1.1 that

$$\max\{\|x(1 - e_\beta)\|_{E(\mathcal{M})}, \|x^*(1 - e_\beta)\|_{E(\mathcal{M})}\} \rightarrow_\beta 0.$$

Consequently, there exists a projection $e_0 \in \mathcal{M}$ with finite trace such that

$$\max\{\|x(1 - e_0)\|_{E(\mathcal{M})}^\alpha, \|x^*(1 - e_0)\|_{E(\mathcal{M})}^\alpha\} < \epsilon^\alpha/2.$$

We next observe that by passing to a subsequence and relabelling if necessary, there exists a sequence e_m of projections in $\mathcal{M}$ with $e_m \uparrow_m e_0$ such that, for all $m = 1, 2, \dots$,

$$\|e_0(x_n - x)e_m e_0\|_\infty \rightarrow_n 0,$$

where $\|\cdot\|_\infty$ denotes the usual operator norm on $\mathcal{M}$. While the proof of this assertion is a standard argument, we include the details for the sake of completeness. In fact, since $e_0(x_n - x)e_0 \rightarrow 0$ for the measure topology given by $(e_0 \mathcal{M} e_0, \tau(e_0 \cdot e_0))$, it follows, by passing to a subsequence if necessary, that there exist projections $e'_k \leq e_0$, $k = 1, 2, \dots$ such that

$$\tau(e_0 - e'_k) < 2^{-k} \quad \text{and} \quad \|e_0(x_k - x)e'_k e_0\|_\infty < 2^{-k}.$$

For $m = 1, 2, \dots$, define $e_m = \inf_{k \geq m} e'_k$. Observe that

$$e_0 - e_m = e_0 - \inf_{k \geq m} e'_k = \sup_{k \geq m} (e_0 - e'_k),$$

and consequently

$$\tau(e_0 - e_m) \leq \sum_{k=m}^{\infty} \tau(e_0 - e'_k) \leq 2^{-m+1},$$

for all $m = 1, 2, \dots$. This implies that $e_m \uparrow_m e_0$. Finally, if $n \geq m$, then

$$\|e_0(x_n - x)e_m e_0\|_{\infty} = \|e_0(x_n - x)e'_n e_m e_0\|_{\infty} \leq \|e_0(x_n - x)e'_n e_0\|_{\infty} \leq 2^{-n},$$

and the assertion follows. Order-continuity of the norm on E again implies that there exists a natural number r such that

$$\max\{\|x(e_0 - e_m)\|_{E(\mathcal{M})}^{\alpha}, \|x^*(e_0 - e_m)\|_{E(\mathcal{M})}^{\alpha}\} < \epsilon^{\alpha}/2, \quad m \geq r.$$

Consequently,

$$\|x(1 - e_r)\|_{E(\mathcal{M})}^{\alpha} \leq \|x(1 - e_0)\|_{E(\mathcal{M})}^{\alpha} + \|x(e_0 - e_r)\|_{E(\mathcal{M})}^{\alpha} < \epsilon^{\alpha},$$

and similarly $\|x^*(1 - e_r)\|_{E(\mathcal{M})} < \epsilon$. We now observe that

$$\|e_r(x_n - x)e_r\|_{\infty} = \|e_0 e_r(x_n - x)e_r e_0\|_{\infty} \leq \|e_0(x_n - x)e_r e_0\|_{\infty}$$

and so

$$\|e_r(x_n - x)e_r\|_{E(\mathcal{M})} \leq \|e_r(x_n - x)e_r\|_{\infty} \|e_0\|_{E(\mathcal{M})}$$

and this suffices to complete the proof of the proposition.

Lemma 2.2 *Let $0 \leq x \in \widetilde{\mathcal{M}}$, let e be a projection in $\mathcal{M}$ with $\tau(e) < \infty$ and let $f = 1 - e$.*

If $0 < \gamma < \infty$, then for all $0 \leq a_1, a_2 \in \mathbf{R}$,

$$\int_0^{a_1} \mu_t^{\gamma}(exe)dt + \int_0^{a_2} \mu_t^{\gamma}(fxf)dt \leq \int_0^{a_1+a_2} \mu_t^{\gamma}(exe + fxf)dt.$$

Proof We begin by observing that, for all $s > 0$,

$$\chi_{(s,\infty)}(|exe + fxf|) = \chi_{(s,\infty)}(|exe|) + \chi_{(s,\infty)}(|fxf|)$$

and so

$$\lambda_s(exe + fxf) = \lambda_s(exe) + \lambda_s(fxf)$$

for all $s > 0$. Using the fact that the functions λ, μ are essential inverses, it follows that

$$\int_0^u \mu_t^\gamma(z) dt = \int_0^\infty u \wedge \lambda_{\frac{1}{s\gamma}}(z) ds$$

for all $z \in \widetilde{\mathcal{M}}$, and for all $0 \leq u \in \mathbf{R}$. Consequently,

$$\begin{aligned} \int_0^{a_1} \mu_t^\gamma(exe) dt + \int_0^{a_2} \mu_t^\gamma(fxf) dt &= \int_0^\infty \left(a_1 \wedge \lambda_{\frac{1}{s\gamma}}(exe) + a_2 \wedge \lambda_{\frac{1}{s\gamma}}(fxf) \right) ds \\ &\leq \int_0^\infty (a_1 + a_2) \wedge \left(\lambda_{\frac{1}{s\gamma}}(exe) + \lambda_{\frac{1}{s\gamma}}(fxf) \right) ds \\ &= \int_0^\infty (a_1 + a_2) \wedge \lambda_{\frac{1}{s\gamma}}(exe + fxf) ds \\ &= \int_0^{a_1+a_2} \mu_t^\gamma(exe + fxf) dt. \end{aligned}$$

We remark that the preceding proof is a variant of the argument of [Fr] Lemma 16.

Lemma 2.3 *Let $x \in \widetilde{\mathcal{M}}$, let e be a projection in $\mathcal{M}$ with $\tau(e) < \infty$ and let $f = 1 - e$. If $0 < \gamma < \infty$, then*

$$\mu^\gamma(e|x|e) + \mu_{(\cdot) - \tau(e)}^\gamma(f|x|f) \prec\prec \mu^\gamma(e|x|e + f|x|f).$$

Proof It suffices to show that if X is a Lebesgue measurable subset of $\mathbf{R}^+$, then

$$\int_X \left[\mu_t^\gamma(e|x|e) + \mu_{t-\tau(e)}^\gamma(f|x|f) \right] dt \leq \int_0^{|X|} \mu_t^\gamma(e|x|e + f|x|f) dt,$$

where $|X|$ denotes the Lebesgue measure of X . Now observe that

$$\begin{aligned} \int_X \left[\mu_t^\gamma(e|x|e) + \mu_{t-\tau(e)}^\gamma(f|x|f) \right] dt &= \int_{X \cap [0, \tau(e)]} \mu_t^\gamma(e|x|e) dt + \int_{X \cap [\tau(e), \infty)} \mu_{t-\tau(e)}^\gamma(f|x|f) dt \\ &\leq \int_0^{|X \cap [0, \tau(e)]|} \mu_t^\gamma(e|x|e) dt + \int_0^{|X \cap [\tau(e), \infty)|} \mu_t^\gamma(f|x|f) dt. \end{aligned}$$

By observing that

$$|X \cap [0, \tau(e)]| + |X \cap [\tau(e), \infty)| = |X|$$

the desired assertion now follows from Lemma 2.2 preceding.

We shall need the following submajorization inequality, which is essentially well known [Si], [GK]. For sake of completeness, we include a proof which is a simple modification of an argument of J. Arazy [Si].

Lemma 2.4 *Let $0 \leq x \in \widetilde{\mathcal{M}}$, let e be a projection in $\mathcal{M}$ and let $f = 1 - e$. If $1 \leq \gamma < \infty$, then*

$$\mu^\gamma(xe + fxf) \prec\prec \mu^\gamma(x).$$

Proof Set $v = e - f$; note that $v \in \mathcal{M}$ is unitary, and that

$$xe + fxf = \frac{1}{2}(x + vxv^*).$$

It follows that

$$\mu(xe + fxf) = \mu\left(\frac{1}{2}(x + vxv^*)\right) \prec\prec \frac{1}{2}(\mu(x) + \mu(vxv^*)) = \mu(x).$$

This shows that the assertion of the Lemma is true if $\gamma = 1$. The case where $\gamma > 1$ follows from the case that $\gamma = 1$, and the fact that the function $(\cdot)^\gamma$ is convex if $\gamma > 1$.

It is not difficult to see that the assertion of Lemma 2.5 fails if $0 < \gamma < 1$, by taking $\mathcal{M} = \mathcal{L}(\mathbf{C}^2)$,

$$x = \begin{pmatrix} 5 & 4 \\ 4 & 5 \end{pmatrix},$$

and by letting e, f be the natural coordinate projections. We omit the details.

We now come to the principal estimate needed for our main result. The inequality given below is a refinement of a similar estimate for trace class operators given by Arazy [Ar] and Lennard [Le1].

Proposition 2.5 *Let E be α -convex with $M^{(\alpha)}(E) = 1$ for some $0 < \alpha \leq 1$ and suppose that E satisfies a lower q -estimate with constant 1 for some finite $q \geq \alpha$. If $k = 2q/\alpha$, then for all $y \in E(\mathcal{M})$ and all projections $e, f \in \mathcal{M}$ with $e + f = 1$ and $\tau(e) < \infty$, it follows that*

$$\|eye\|_{E(\mathcal{M})}^k + \|eyf\|_{E(\mathcal{M})}^k + \|fye\|_{E(\mathcal{M})}^k + \|fyf\|_{E(\mathcal{M})}^k \leq \|y\|_{E(\mathcal{M})}^k.$$

Proof The stated assumptions on E imply that $E^{1/\alpha}$ is a Banach lattice which satisfies a lower (q/α) -estimate, with constant 1. Let $0 \leq w \in E^{(1/\alpha)}(\mathcal{M})$ and let e, f be as stated. Using the lower estimate, it follows that

$$\begin{aligned} \|ewe\|_{E^{(1/\alpha)}(\mathcal{M})}^{q/\alpha} + \|fwf\|_{E^{(1/\alpha)}(\mathcal{M})}^{q/\alpha} &= \|\mu(ewe)\|_{E^{(1/\alpha)}}^{q/\alpha} + \|\mu_{(\cdot) - \tau(e)}(fwf)\|_{E^{(1/\alpha)}}^{q/\alpha} \\ &\leq \|\mu(ewe) + \mu_{(\cdot) - \tau(e)}(fwf)\|_{E^{(1/\alpha)}}^{q/\alpha} \\ &\leq \|\mu(ewe + fwf)\|_{E^{(1/\alpha)}}^{q/\alpha} \\ &\leq \|\mu(w)\|_{E^{(1/\alpha)}}^{q/\alpha} = \|w\|_{E^{(1/\alpha)}(\mathcal{M})}^{q/\alpha}, \end{aligned}$$

where the last two inequalities follow from successive application of Lemmas 2.3, 2.4. It now follows that for all $z \in E^{(2/\alpha)}(\mathcal{M})$ that

$$\|ez\|_{E^{(2/\alpha)}(\mathcal{M})}^k + \|fz\|_{E^{(2/\alpha)}(\mathcal{M})}^k \leq \|z\|_{E^{(2/\alpha)}(\mathcal{M})}^k. \quad (*)$$

In fact,

$$\begin{aligned} \|ez\|_{E^{(2/\alpha)}(\mathcal{M})}^k + \|fz\|_{E^{(2/\alpha)}(\mathcal{M})}^k &= \|(ezz^*e)^{1/2}\|_{E^{(2/\alpha)}(\mathcal{M})}^k + \|(fzz^*f)^{1/2}\|_{E^{(2/\alpha)}(\mathcal{M})}^k \\ &= \|ezz^*e\|_{E^{(1/\alpha)}(\mathcal{M})}^{k/2} + \|fzz^*f\|_{E^{(1/\alpha)}(\mathcal{M})}^{k/2} \\ &\leq \|zz^*\|_{E^{(1/\alpha)}(\mathcal{M})}^{k/2} = \|z\|_{E^{(2/\alpha)}(\mathcal{M})}^k. \end{aligned}$$

Set $\beta = 2/\alpha$ and choose γ , $1 < \gamma \leq \infty$, such that $2/\beta + 1/\gamma = 1$, which is possible since $\beta \geq 2$. Suppose now that $y \in E(\mathcal{M})$ and set

$$a = u|y|^{1/\beta}, \quad b = |y|^{1/\gamma}, \quad c = |y|^{1/\beta},$$

where $y = u|y|$ is the polar decomposition of y . It follows that $a, c \in E^{(\beta)}(\mathcal{M})$ and $b \in E^{(\gamma)}(\mathcal{M})$ (if $\gamma = \infty$, we take b to be 1 and $E^{(\gamma)}(\mathcal{M})$ to be $\mathcal{M}$). Note that

$$\|a\|_{E^{(\beta)}(\mathcal{M})}^\beta = \|b\|_{E^{(\gamma)}(\mathcal{M})}^\gamma = \|c\|_{E^{(\beta)}(\mathcal{M})}^\beta = \|y\|_{E(\mathcal{M})}. \quad (**)$$

We apply the Hölder-type inequality (1.3) to obtain

$$\begin{aligned}\|eye\|_{E(\mathcal{M})} &= \|eabce\|_{E(\mathcal{M})} \\ &\leq \|ea\|_{E^{(\beta)}(\mathcal{M})} \|b\|_{E^{(\gamma)}(\mathcal{M})} \|ce\|_{E^{(\beta)}(\mathcal{M})}.\end{aligned}$$

Similarly,

$$\begin{aligned}\|eyf\|_{E(\mathcal{M})} &\leq \|ea\|_{E^{(\beta)}(\mathcal{M})} \|b\|_{E^{(\gamma)}(\mathcal{M})} \|cf\|_{E^{(\beta)}(\mathcal{M})}, \\ \|fye\|_{E(\mathcal{M})} &\leq \|fa\|_{E^{(\beta)}(\mathcal{M})} \|b\|_{E^{(\gamma)}(\mathcal{M})} \|ce\|_{E^{(\beta)}(\mathcal{M})}, \\ \|fyf\|_{E(\mathcal{M})} &\leq \|fa\|_{E^{(\beta)}(\mathcal{M})} \|b\|_{E^{(\gamma)}(\mathcal{M})} \|cf\|_{E^{(\beta)}(\mathcal{M})}.\end{aligned}$$

Consequently, using first (*) and then (**) we obtain that

$$\begin{aligned}&\|eye\|_{E(\mathcal{M})}^k + \|eyf\|_{E(\mathcal{M})}^k + \|fye\|_{E(\mathcal{M})}^k + \|fyf\|_{E(\mathcal{M})}^k \\ &\leq \left(\|ea\|_{E^{(\beta)}(\mathcal{M})}^k + \|fa\|_{E^{(\beta)}(\mathcal{M})}^k \right) \|b\|_{E^{(\gamma)}(\mathcal{M})}^k \left(\|cf\|_{E^{(\beta)}(\mathcal{M})}^k + \|ce\|_{E^{(\beta)}(\mathcal{M})}^k \right) \\ &\leq \|a\|_{E^{(\beta)}(\mathcal{M})}^k \|b\|_{E^{(\gamma)}(\mathcal{M})}^k \|c\|_{E^{(\beta)}(\mathcal{M})}^k \\ &= \|y\|_{E(\mathcal{M})}^{k(2/\beta+1/\gamma)} \\ &= \|y\|_{E(\mathcal{M})}^k,\end{aligned}$$

and the proof of the proposition is complete.

Before proceeding, it is convenient to introduce the following terminology.

Definition 2.6 Let $(X, \|\cdot\|_X)$ be an α -normed quasi-Banach space, for some $0 < \alpha \leq 1$, with unit ball $\mathbf{B}_X$, and let $\mathcal{T}$ be a topological vector space topology on X that is weaker than the quasi-norm topology. The space $(X, \|\cdot\|_X)$ is said to have the uniform Kadec-Klee-Huff property with respect to $\mathcal{T}$ (denoted $UKKH(\mathcal{T})$) if and only if for every $\epsilon > 0$ there exists $\delta = \delta(\epsilon) \in (0, 1)$ such that whenever $\{x_n\}$ is a sequence in $\mathbf{B}_X$ with $x_n \rightarrow_n x \in X$ with respect to $\mathcal{T}$ and $\|x\|_X > 1 - \delta$, it follows that for some positive integer N ,

$$\sup_{n, m \geq N} \|x_n - x_m\|_X \leq \epsilon.$$

Let us immediately note that the conclusion in the statement of the preceding definition is stronger than the usual formulation of this property given by several authors (cf

[H], [Le1,2], [DL1,2]) in which sup is replaced by inf. However, a minor modification of the argument of [BDDL] Proposition 1.2 shows that our (apparently) stronger formulation is, in fact, equivalent to that already in the literature.

We may now state the main result of this paper.

Theorem 2.7 *Let E be a symmetric quasi-Banach functions space on the positive half-line $\mathbf{R}^+$. If E is α -convex with $M^{(\alpha)}(E) = 1$ for some $0 < \alpha \leq 1$ and if E satisfies a lower- q estimate with constant 1 for some finite $q \geq \alpha$, then $(E(\mathcal{M}), \|\cdot\|_{E(\mathcal{M})})$ has the uniform Kadec-Klee-Huff property with respect to local convergence in measure.*

Proof Suppose that $\delta \in (0, 1)$ and that $\{x_n\}_{n=1}^\infty$ is a sequence in the unit ball of $E(\mathcal{M})$ satisfying $x_n \rightarrow_n x$ locally in measure for some $x \in E(\mathcal{M})$ with $\|x\|_{E(\mathcal{M})} > 1 - \delta$. By [BDDL], Proposition 1.2, it suffices to show that for some continuous, invertible function $\epsilon = \epsilon(\delta)$ mapping $(0, \delta_0)$ into $(0, \infty)$, where $\delta_0 > 0$ and $\epsilon(\delta) \rightarrow 0$ as $\delta \rightarrow 0$, we have

$$\inf_{n \neq m} \|x_n - x_m\|_{E(\mathcal{M})} < \epsilon.$$

By passing to a subsequence and relabelling, if necessary, and appealing to Proposition 2.1, we may assume that there exists a projection $e \in \mathcal{M}$ with $\tau(e) < \infty$ such that, setting $f = 1 - e$,

$$\|e(x_n - x)e\|_{E(\mathcal{M})} \rightarrow_n 0 \tag{i}$$

and

$$\max \{ \|xf\|_{E(\mathcal{M})}, \|x^*f\|_{E(\mathcal{M})} \} < \left[\frac{1}{3} \left(\|x\|_{E(\mathcal{M})}^\alpha - (1 - \delta)^\alpha \right) \right]^{1/\alpha}. \tag{ii}$$

Since $\|\cdot\|_{E(\mathcal{M})}$ is an α -norm, we obtain that

$$\|x_n - x_m\|_{E(\mathcal{M})}^\alpha \leq \|e(x_n - x_m)e\|_{E(\mathcal{M})}^\alpha + R(x_n) + R(x_m),$$

where

$$R(y) = \|eyf\|_{E(\mathcal{M})}^\alpha + \|fye\|_{E(\mathcal{M})}^\alpha + \|fyf\|_{E(\mathcal{M})}^\alpha,$$

for all $y \in E(\mathcal{M})$. Setting $k = 2q/\alpha$, $s = k/\alpha$ and $s' = s/(s-1)$, and applying Proposition 2.5, we obtain, for each $y \in \mathcal{M}$,

$$\begin{aligned} R(y) &\leq 3^{1/s'} \left(\|eyf\|_{E(\mathcal{M})}^k + \|fye\|_{E(\mathcal{M})}^k + \|fyf\|_{E(\mathcal{M})}^k \right)^{1/s} \\ &\leq 3^{1/s'} \left(\|y\|_{E(\mathcal{M})}^k - \|eye\|_{E(\mathcal{M})}^k \right)^{1/s}. \end{aligned}$$

Using (ii), it now follows that

$$\begin{aligned} \|exe\|_{E(\mathcal{M})}^\alpha &\geq \|x\|_{E(\mathcal{M})}^\alpha - \|exf\|_{E(\mathcal{M})}^\alpha - \|fxe\|_{E(\mathcal{M})}^\alpha - \|fxf\|_{E(\mathcal{M})}^\alpha \\ &\geq \|x\|_{E(\mathcal{M})}^\alpha - 3 \max \left\{ \|xf\|_{E(\mathcal{M})}^\alpha, \|x^*f\|_{E(\mathcal{M})}^\alpha \right\} \\ &> \|x\|_{E(\mathcal{M})}^\alpha - \left(\|x\|_{E(\mathcal{M})}^\alpha - (1-\delta)^\alpha \right) \\ &= (1-\delta)^\alpha, \end{aligned}$$

so that $\|exe\|_{E(\mathcal{M})} > 1-\delta$. From (i) and the uniform continuity of the quasi-norm $\|\cdot\|_{E(\mathcal{M})}$ on the closed unit ball of $E(\mathcal{M})$, it follows that

$$\|ex_n e\|_{E(\mathcal{M})} \rightarrow_n \|exe\|_{E(\mathcal{M})} ;$$

and so there exists a natural number n_0 such that

$$\|ex_n e\|_{E(\mathcal{M})} > 1-\delta, \quad n \geq n_0.$$

Consequently, for all $n, m \geq n_0$, it follows that

$$\begin{aligned} \|x_n - x_m\|_{E(\mathcal{M})}^\alpha &\leq \|e(x_n - x_m)e\|_{E(\mathcal{M})}^\alpha + 3^{1/s'} \left(\|x_n\|_{E(\mathcal{M})}^k - \|ex_n e\|_{E(\mathcal{M})}^k \right)^{1/s} \\ &\quad + 3^{1/s'} \left(\|x_m\|_{E(\mathcal{M})}^k - \|ex_m e\|_{E(\mathcal{M})}^k \right)^{1/s} \\ &\leq \|e(x_n - x_m)e\|_{E(\mathcal{M})}^\alpha + 2 \cdot 3^{1/s'} (1 - (1-\delta)^k)^{1/s}. \end{aligned}$$

We obtain that

$$\inf_{n \neq m} \|x_n - x_m\|_{E(\mathcal{M})} \leq Q(1 - (1-\delta)^k)^{1/k},$$

where $Q = 2^{1/\alpha} 3^{1/(\alpha s')}$. Thus, given $\epsilon \in (0, Q)$, we may take $\delta(\epsilon)$ to be given by the formula

$$\delta(\epsilon) = 1 - \left(1 - (\epsilon/Q)^k \right)^{1/k} ;$$

and by this the proof of the theorem is complete.

In order to state some explicit consequences of the preceding theorem, let us recall (cf. [LT] p.142) that, for $0 < p \leq q < \infty$, the Lorentz space $L_{q,p}(\mathbf{R}^+)$ consists of all Lebesgue measurable functions f on $\mathbf{R}^+$ for which

$$\|f\|_{q,p} = \left(\int_0^\infty \mu_t^p(f) dt^{p/q} \right)^{1/p} < \infty$$

It is shown in [Cr] Proposition 3.2 that $L_{q,p}(\mathbf{R}^+)$ is p -convex with $M^{(p)}(L_{q,p}(\mathbf{R}^+)) = 1$ and satisfies a lower q -estimate with constant 1, in the case that $p \geq 1$; however, it is not difficult to see that this restriction is unnecessary and further that $L_{q,p}(\mathbf{R}^+)$ is a p -Banach space when $0 < p < 1$. We may now state the following consequence of Theorem 2.7.

Corollary 2.8. *If $0 < p \leq q < \infty$ then the Lorentz space $L_{q,p}(\mathcal{M})$ has the uniform Kadec-Klee-Huff property with respect to local convergence in measure.*

The corollary appears to be new even in the case that $\mathcal{M}$ is commutative, and in this setting, the corollary extends earlier results in [Le2,3]. As has been noted earlier, if $\mathcal{M}$ is the space $\mathcal{L}(H)$ of all bounded linear operators on H and τ is the standard trace, then convergence locally in measure is precisely convergence for the weak operator topology. In this case, the corollary implies that the Schatten p -classes $\mathcal{C}_p$, $0 < p < \infty$, have the uniform Kadec-Klee-Huff property for the weak operator topology. For the Schatten class $\mathcal{C}_1$, this is again due to Lennard [Le1]. Of course, if $1 < p < \infty$, then the result for $\mathcal{C}_p$ may be derived directly by using uniform convexity and Clarkson's inequalities.

We remark further, that if $\mathcal{M}$ is commutative and the underlying measure space is σ -finite, or if $\mathcal{M}$ is the space $\mathcal{L}(H)$ of all bounded linear operators on a separable Hilbert space H with the usual trace, or if $\tau(1) < \infty$, then the topology of local convergence in measure is a metric topology. Accordingly, when combined with [Le2] Corollary 3.3 and Theorem 4.2, our Theorem 2.7 immediately yields results concerning normal structure and

fixed point properties in each of the above cases. We leave precise formulations to the interested reader. We state, however a further consequence of Theorem 2.7 and [BDDL] Theorem 3.1, which seems to be of independent interest.

Corollary 2.9 *If (Ω, Σ, μ) is a measure space, if $1 \leq p < \infty$, and if E satisfies the conditions of Theorem 2.7 with $\alpha = 1$, then the Lebesgue-Bochner space $L^p(\mu, E(\mathcal{M}))$ has the following property. For each $\epsilon > 0$, there exists $\delta \in (0, 1)$ such that whenever $\{f_n\}$ is a sequence in $\mathbf{B}_{L^p(E(\mathcal{M}))}$ and $f \in L^p(E(\mathcal{M}))$ with $f_n(\omega) \rightarrow_n f(\omega)$ locally in measure for almost all $\omega \in \Omega$ and $\|f\|_{L^p(E(\mathcal{M}))} > 1 - \delta$, then for some positive integer N*

$$\sup_{n, m \geq N} \|f_n - f_m\|_{L^p(E(\mathcal{M}))} \leq \epsilon .$$

We remark that, as noted in [BDDL], the above result is still true if $0 < p < 1$. Moreover, the result still holds if $0 < \alpha < 1$.

Theorem 2.10 *Let E be a symmetric Banach function space on $\mathbf{R}^+$. If E has $UKKH$ (lcm), then E satisfies a lower q -estimate for some finite q .*

Proof If E fails to satisfy a lower q -estimate for some finite q , then it follows from [LT] 1.f.12 that for every $\epsilon > 0$ and $n \in \mathbf{N}$, there exists a (finite) sequence $\{x_i\}_{i=1}^n$ of mutually disjoint elements of E such that

$$\max_{1 \leq i \leq n} |a_i| \leq \left\| \sum_{i=1}^n a_i x_i \right\|_E \leq (1 + \epsilon) \max_{1 \leq i \leq n} |a_i|,$$

for all choices of scalars $\{a_i\}_{i=1}^n$.

Let $\delta > 0$ be fixed and choose disjoint elements $x, y \in E$ such that

$$\max\{|a|, |b|\} \leq \|ax + by\|_E \leq (1 + \delta) \max\{|a|, |b|\}$$

for every choice of scalars a, b . Observe that

$$1 \leq \|x\|_E, \|y\|_E, \|x + y\|_E \leq 1 + \delta.$$

Set

$$x' = \|x + y\|_E^{-1}x, \quad y' = \|x + y\|_E^{-1}y$$

so that $\|x' + y'\|_E = 1$ and

$$(1 + \delta)^{-1} \leq \|x'\|_E, \|y'\|_E \leq 1 + \delta.$$

For each $n = 0, 1, 2, \dots$, set $I_n = [2n + 1, 2n + 2)$ and let $T = \cup_{n=0}^{\infty} I_n$. Let $\{T_n\}_{n=1}^{\infty}$ be a measurable partition of T into a mutually disjoint sequence of subsets of T with $|T_n| = \infty$, $n = 1, 2, \dots$. Let $\phi : \mathbf{R}^+ \rightarrow \mathbf{R}^+ \setminus T$ and $\phi_n : \mathbf{R}^+ \rightarrow T_n$, $n = 1, 2, \dots$ be measure preserving bijections. Set

$$x'' = x' \circ \phi^{-1}, \quad y_n = y' \circ \phi_n^{-1}, \quad z_n = x'' + y_n$$

for $n = 1, 2, \dots$. It follows that

$$\|z_n\|_E = \|x'' + y_n\|_E = \|x' + y'\|_E = 1,$$

and

$$\begin{aligned} \|z_n - z_m\|_E &= \|x'' + y_n - (x'' + y_m)\|_E = \|y_n - y_m\|_E \\ &\geq \|y_n\|_E = \|y'\|_E \geq (1 + \delta)^{-1} \end{aligned}$$

for all $n, m = 1, 2, \dots$ with $n \neq m$. Since $y_n \rightarrow 0$ locally in measure, it follows that $z_n \rightarrow x''$ locally in measure. However, since $\|x''\|_E \geq (1 + \delta)^{-1}$ and since $\delta > 0$ is arbitrary, it follows that E does not have $UKKH(lcm)$.

If E is a symmetric Banach function space on $\mathbf{R}^+$, and if E satisfies a lower q -estimate for some $1 < q < \infty$, then it follows from [LT] 1.f.7 that E is r -concave for every $r > q$. For each such $r > q$, the space E may be renormed equivalently so that E , endowed with the new norm and the same order is a symmetric Banach function space on $\mathbf{R}^+$ which is r -concave with r -concavity constant 1. See, for example, [LT] 1.d.8, 1.f.11. We obtain therefore the following consequences.

Corollary 2.11 *If E is a symmetric Banach function space on $\mathbf{R}^+$, then E satisfies a lower q -estimate for some $1 < q < \infty$ if and only if there is an equivalent symmetric norm $\|\cdot\|_0$ on E such that $(E, \|\cdot\|_0)$ has $UKKH(lcm)$.*

Corollary 2.12 *If E is a symmetric Banach function space on $\mathbf{R}^+$ and if E has $UKKH(lcm)$, then there is an equivalent norm $\|\cdot\|_0$ on $E(\mathcal{M})$ such that $(E(\mathcal{M}), \|\cdot\|_0)$ has $UKKH(lcm)$.*

3. Final remarks and related results There are several variants of the commutative specialization of Theorem 2.7. In this section, we formulate one such variant in the setting of Banach lattices. We begin with the following simple characterization of Banach lattices which are q -concave, for some finite q .

Proposition 3.1 *If E is a Banach lattice, then the following statements are equivalent.*

- (i) *E satisfies a lower q -estimate for some $q < \infty$.*
- (ii) *E admits an equivalent norm $\|\cdot\|_0$ such that E equipped with $\|\cdot\|_0$ and the same order is a Banach lattice with the following property: For every $c > 0$, there exists $0 < \delta = \delta(c)$ such that, whenever $x, y, z \in E^+$ satisfy*

$$x = y + z, \quad y \wedge z = 0, \quad \|y\|_0 = 1, \quad \|z\|_0 \geq c,$$

it follows that $\|x\|_0 \geq 1 + \delta(c)$.

That (i) implies (ii) is a simple consequence of [LT] 1.f.7, 1.d.8. The reverse implication follows from [LT] 1.f.12, and the details are omitted.

A Banach lattice E which satisfies the stated condition (ii) in Proposition 3.1 preceding in its given norm is said to have property (C). We remark that it follows from Proposition 3.1 that if a Banach lattice E has property (C), then E necessarily has order continuous norm.

Definition 3.2 *Let E be a Banach lattice. The sequence $\{x_n\} \subseteq E$ is said to converge locally to $x \in E$ if and only if for every band projection Q on E there exists a band projection P with $0 \neq P \leq Q$ and a subsequence $\{y_n\} \subseteq \{x_n\}$ such that $\|P(y_n - x)\| \rightarrow_n 0$.*

Let us mention immediately that if E is any Banach function space with order continuous norm on some σ -finite measure space, and if the sequence $\{x_n\} \subseteq E$ converges to $x \in E$ in measure on every subset of finite measure, then it follows from Egorov's theorem that the sequence $\{x_n\}$ converges locally to x .

The theorem which follows, and its proof, is an adaptation of a similar result due to van Dulst and de Valk [DV], Proposition 3, in the setting of Banach spaces with a Schauder basis. We include the details of proof for the sake of completeness.

Theorem 3.3 *Let E be a Banach lattice. If E has property (C), then E has the following property. For every $\epsilon > 0$, there exists $\delta = \delta(\epsilon) \in (0, 1)$ such that whenever $\{x_n\}$ is a sequence in the unit ball of E which converges locally to $x \in E$, and which satisfies $\|x_n - x_m\| > \epsilon, n \neq m$, it follows that $\|x\| \leq 1 - \delta$.*

Proof Let $\epsilon > 0$ be given and assume that there is no $\delta > 0$ satisfying the assertion of the theorem for this ϵ . For each $c > 0$, let $\delta(c) > 0$ be the largest δ which satisfies the assertion of property (C), and let $0 < \alpha < 1$ satisfy $\alpha(1 + \delta(\epsilon/2)) > 1$. We assume that there exists a sequence $\{x_n\}$ in the unit ball of E which satisfies $\|x_n - x_m\| > \epsilon, n \neq m$, and which converges locally to some $x \in E$ with $\|x\| > \alpha$. Observe that local convergence of the sequence $\{x_n\}$ to x together with order continuity of the norm on E implies that there exists a band projection P and a subsequence $\{y_n\} \subseteq \{x_n\}$ such that $\|Px\| > \alpha$ and $\|P(y_n - x)\| \rightarrow_n 0$. Consequently there exists a natural number n_0 such that

$$\|Py_n\| > \alpha, \quad \|(I - P)(y_n - y_m)\| > \epsilon$$

for all natural numbers $n, m \geq n_0, n \neq m$. For at least one of the choices $n = n_0, n_0 + 1$,

it follows that

$$\alpha < \|Py_n\| \leq 1, \quad \|(I - P)y_n\| > \epsilon/2.$$

With this choice of index, set

$$w = \frac{y_n}{\|Py_n\|}, \quad y = P\left(\frac{y_n}{\|Py_n\|}\right), \quad z = (I - P)\left(\frac{y_n}{\|Py_n\|}\right),$$

and observe that

$$|w| = |y| + |z|, \quad |y| \wedge |z| = 0, \quad \| |y| \| = 1, \quad \| |z| \| \geq \epsilon/2.$$

Property (C) now implies that $\| |w| \| \geq 1 + \delta(\epsilon/2)$ so that

$$\|y_n\| = \| |y_n| \| > \alpha(1 + \delta(\epsilon/2)) > 1,$$

and this is a contradiction.

References

- [Ar] J. Arazy, *More on convergence in unitary matrix spaces*, Proc. Amer. Math. Soc. **83** (1981), 44-48.
- [BDDL] M. Besbes, S.J. Dilworth, P.N. Dowling, and C.J. Lennard, *New Convexity and fixed point properties in Hardy and Lebesgue-Bochner spaces*, J. Funct. Anal. (to appear)
- [CDLT] N.L. Carothers, S.J. Dilworth, C.J. Lennard and D.A. Trautman, *A fixed point property for the Lorentz space $L_{p,1}(\mu)$* , Indiana Univ. Math. J. **40** (1991) 345-352.
- [Cr] J. Creekmore, *Type and cotype in Lorentz $L_{p,q}$ spaces*, Indag. Math., **43** (1981) 145-152.
- [CS] V.I. Chi'lin and F.A. Sukochev, *Convergence in measure in admissible non-commutative symmetric spaces*, Izv. Vysš Učeb. Zaved **9**(1990) 63-90 (Russian).
- [CKS] V.I. Chi'lin, A.V. Krygin and P.A. Sukochev, *Uniform and local uniform convexity*

of symmetric spaces of measurable operators,, Dep VINITI N5620-B90(1990) 24pp. (Russian).

[DH] S.J. Dilworth and Y.P. Hsu, *The uniform Kadec-Klee property in the trace ideals $\mathcal{C}_E$* , preprint 1993.

[DDP1] P.G. Dodds, T.K. Dodds and B. de Pagter, *Non-commutative Banach function spaces*, Math. Z. **201** (1989), 583-597.

[DDP2] P.G. Dodds, T.K. Dodds and B. de Pagter, *Non-commutative Köthe duality*, Trans. Amer. Math. Soc. (to appear).

[DL1] P.N. Dowling and C.J. Lennard, *Kadec-Klee properties of vector-valued Hardy spaces*, Math. Proc. Camb. Phil. Soc. **111** (1992) 535-544.

[DL2] P.N. Dowling and C.J. Lennard, *On uniformly H -convex complex quasi-Banach spaces*, Bull. des Sci. Math. (to appear)

[DV] D.van Dulst and V.de Valk, *(KK)-properties, normal structure and fixed points of nonexpansive mappings in Orlicz sequence spaces*, Can. J. Math. **38** (1986) 728-750.

[Fa] T. Fack, *Sur la notion de valeur caractéristique*, J. Operator Theory **7** (1982), 307-333.

[FK] T. Fack and H. Kosaki, *Generalized s -numbers of τ -measurable operators*, Pacific J. Math. **123** (1986), 269-300.

[Fr] D.H. Fremlin, *Stable subspaces of $L^1 + L^\infty$* , Math. Proc. Camb. Philos. Soc. **64** (1968) 625-643.

[GK] I.C. Gohberg and M.G. Krein, *Introduction to the theory of non-selfadjoint operators*, Translations of Mathematical Monographs, vol.18, AMS (1969).

[H] R. Huff, *Banach spaces which are nearly uniformly convex*, Rocky Mountain J. Math. **10** (1980) 743-749.

[KPS] S.G. Krein, Ju.I. Petunin and E.M. Semenov, *Interpolation of linear operators*, Translations of Mathematical Monographs, vol.54, AMS (1982).

[Le1] C. Lennard, *$\mathcal{C}_1$ is uniformly Kadec-Klee*, Proc. Amer. Math. Soc. **109** (1990) 71-77.

[Le2] C. Lennard, *A new convexity property that implies a fixed point property for L_1* ,

Studia Math. **100** (1991) 95-108.

[Le3] C. Lennard, *Operators and geometry of Banach spaces*, PhD Thesis, Kent State University, 1988.

[LT] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces II, Function Spaces*, Springer-Verlag, 1979.

[M] C.A. McCarthy, c_p , Israel J. Math. **5** (1967), 249-271.

[Ne] E. Nelson, *Notes on non-commutative integration*, J. Funct. Anal. **15** (1974), 103-116.

[Ov] V.I. Ovcinnikov, *s-numbers of measurable operators*, Funktsional'nyi Analiz i Ego Prilozheniya **4** (1970) 78-85 (Russian).

[Si] B. Simon, *Convergence in trace ideals*, Proc. Amer. Math. Soc. **83**(1981), 39-43.

[Su] F.A. Sukochev, *On the uniform Kadec-Klee property*, International Conference dedicated to the 100-th birthday of S.Banach, Lwow, 1992.

[Te] M. Terp, *L^p -spaces associated with von Neumann algebras*, Notes, Copenhagen University (1981).

[X] Q. Xu, *Analytic functions with values in lattices and symmetric spaces of measurable operators*, Math. Proc. Camb. Phil. Soc. **109** (1991) 541-563.

Addresses

P.G. Dodds

School of Information Sci. & Technology

The Flinders University of South Australia

GPO Box 2100, Adelaide 5001

South Australia

T.K. Dodds

School of Information Sci. & Technology

The Flinders University of South Australia

GPO Box 2100, Adelaide 5001

South Australia

P.N. Dowling

C.J. Lennard

Department of Mathematics and Statistics	Department of Mathematics and Statistics
Miami University	University of Pittsburgh
Oxford, Ohio 45056	Pittsburgh, Pennsylvania 15260

F.A. Sukochev

Department of Mathematics
Tashkent State University
Vuzgorodok, 700095, Tashkent
Uzbekistan